impossible if the sympathetic were inhibitor. In animals such as the cat and guinea-pig reflexes through the sympathetic to the internal generative organs during coitus, leading to simultaneous contraction of the vagina and relaxation of the uterus, may aid in the injection of spermatozoa into the uterine cavity.

Conclusion.

In the rabbit, cat, guinea-pig and rat (and by analogy probably in all animals) the sympathetic nerve supply of the vagina is predominantly motor in quality, whether it be motor or inhibitor to the uterus.

REFERENCES.

1. Langley and Anderson, 'Journ. of Phys.,' vol. 19, p. 122 (1895).
2. Cushny, *ibid.*, vol. 35, p. 1 (1906).
3. Dale, *ibid.*, vol. 34, p. 163 (1906).
4. Gunn and Gunn, 'Journ. Pharmacol. Expt. Ther.,' vol. 5, p. 527 (1914).
5. Waddell, 'Journ. Pharmacol. Expt. Ther.,' vol. 9, p. 411 (1911).
6. Gunn and Davies, 'Journ. of Phys.,' vol. 54 ('Proceedings,' November, 1920).

An Investigation of the Colour Vision of 527 *Students by the Rayleigh Test.*

By Dr. R. A. Houston.

[This paper is published in Series A, No. 716, December, 1922.]